

**IMAGE**

Short-scar pectoralis major flap through a submammary fold incision

Daniel Sattler¹, Maria von Kohout¹, Armin Kraus²

¹Department of Plastic and Aesthetic Surgery, Beta Clinic, Bonn;

²Department of Plastic, Aesthetic and Hand Surgery, Otto von Guericke University, Magdeburg, Germany

Correspondence: Armin Kraus

Department of Plastic, Aesthetic and Hand Surgery, Otto von Guericke University, Leipziger Strasse 44, D-39120 Magdeburg, Germany
Tel: +49-391-67-15599, Fax: +49-391-67-15588
E-mail: armin.kraus@med.ovgu.de

Received: December 22, 2018 • Revised: May 24, 2019 • Accepted: May 29, 2019
pISSN: 2234-6163 • eISSN: 2234-6171
<https://doi.org/10.5999/aps.2018.01543>
Arch Plast Surg 2019;46:486-487

Copyright © 2019 The Korean Society of Plastic and Reconstructive Surgeons
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

We report the case of a 19-year-old female patient who required supraclavicular defect coverage. She suffered from congenital brachial plexus palsy and had undergone 17 prior operations. Due to instability, the clavicle had to be shortened. Wound dehiscence and piercing of the end of the clavicle through the skin occurred (Fig. 1). As the patient desired a maximally safe operation with an optimal aesthetic result, we chose a short-scar pectoralis major flap (PMF). The pectoralis major (PM) muscle appeared clinically unaffected, while the back showed extensive scarring from prior operations. A magnetic resonance imaging scan depicted the thoracoacromial artery. The patient provided written informed consent for



Fig. 1.

Tissue defect with exposure of the clavicle. Soft tissue defect with exposure of the left clavicle after several attempts to perform secondary sutures, with extensive scarring on the back.

surgery and publication. The operation was rated as a category C procedure according to Bernstein and Bampoe [1] (amendment to the technique of an established operation) and therefore exempt from institutional review board approval. Debridement with smoothing of the clavicular end was performed first. An 8-cm submammary incision was made and the PM was identified. After dissection, the flap was flipped cranially, pulled through a tunnel up into the defect and fixed above the clavicular stump (Fig. 2). The postoperative result was aesthetically pleasing (Fig. 3). Defect coverage in the shoulder region by PMFs is not new, and PMFs compete with a vast choice of free fasciocutaneous and musculocutaneous flaps. The complications of PMFs range from breast distortion to complete flap necrosis [2]. The PMF is not known for good cosmesis, usually leaving extensive scars. An inframammary approach was described by Zbar et al. [3], but with an incision leaving a longer scar. This led us to think about a scar-sparing approach, and making an incision of the type used for breast augmentation seemed natural.



Fig. 2.

Defect coverage with the pectoralis major muscle. Soft tissue coverage of the supraclavicular region by a pectoralis major flap.



Fig. 3.

Postoperative view 2 weeks after surgery. Good defect coverage and favorable aesthetic results after flap elevation through the submammary fold incision were obtained.

Notes

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Ethical approval

The study was performed in accordance with the

principles of the Declaration of Helsinki. Written informed consent was obtained.

Patient consent

The patient provided written informed consent for the publication and the use of her images.

Author contribution

Methodology, data curation: Sattler S, von Kohout M, Kraus A. Project administration: Sattler S. Visualization, writing original draft: Kraus A. Review&editing: Sattler S, von Kohout M. Approval of final manuscript: all authors.

ORCID

Daniel Sattler <https://orcid.org/0000-0002-9529-8115>

Maria von Kohout <https://orcid.org/0000-0001-8691-5057>

Armin Kraus <https://orcid.org/0000-0001-6557-2163>

References

1. Bernstein M, Bampoe J. Surgical innovation or surgical evolution: an ethical and practical guide to handling novel neurosurgical procedures. *J Neurosurg* 2004;100:2-7.
2. Kruse AL, Luebbers HT, Obwegeser JA, et al. Evaluation of the pectoralis major flap for reconstructive head and neck surgery. *Head Neck Oncol* 2011;3:12.
3. Zbar RI, Funk GF, McCulloch TM, et al. Pectoralis major myofascial flap: a valuable tool in contemporary head and neck reconstruction. *Head Neck* 1997;19:412-8.